

**Adam Seidl, MD**  
Assistant Professor  
Shoulder/Elbow Surgery  
Department of Orthopedics  
University of Colorado



**Orthopedics**  
UNIVERSITY OF COLORADO

**Highlands Ranch Hospital**  
(720) 516-4090  
**Inverness**  
(303) 694-3333  
**Anschutz Medical Campus**  
(720) 848-1900

## **Rehabilitation Protocol**

### **Non-operative Acromioclavicular Joint Separation**

#### **Phase I: Gradual ROM** (*Injury to 2 weeks*)

- Ice encouraged for the first week at a minimum: should be used 3-4 times per day.
- Sling should be in place for 5-7 days when not performing exercises.
- Initiate exercise program 3 times per day:
  - Immediate elbow, forearm and hand range of motion out of sling
  - Passive and active assistive ER at the side to 30, flexion to 130
- No lifting with involved extremity.
- AVOID scapular ROM exercises.

#### **Phase II: Progress ROM & Function** (*2 to 6 weeks*)

- May discontinue sling.
- Advance active and passive ROM in all planes to tolerance.
- Lifting restriction of 5 pounds with the involved extremity until 4 weeks from injury
- Initiate gentle rotator cuff strengthening; progress to light resistive Theraband exercises.
- Initiate scapular AROM exercises.

#### **Phase III: Full Function** (*>6 weeks*)

- Discontinue all lifting restrictions.
- Advance rotator cuff and scapular stabilizer strengthening.
- Initiate functional progression to sports specific activities at 6-8 weeks.